

Estimates of $m_d - m_u$ and $\langle \bar{d}d \rangle - \langle \bar{u}u \rangle$ from QCD sum rules for D and D^* isospin mass differences

V.L. Eletsky[†],

Theoretical Physics Institute, University of Minnesota
Minneapolis, MN 55455, USA

and

Institute for Theoretical Physics, Bern University
Sidlerstrasse 5, CH-3012 Bern, Switzerland

and

B.L. Ioffe

Institute of Theoretical and Experimental Physics
Moscow 117259, Russia

Abstract

The recent experimental data on $D^+ - D^0$ and $D^{*+} - D^{*0}$ mass differences are used as inputs in the QCD sum rules to obtain new estimates on the mass difference of light quarks and on the difference of their condensates: $m_d - m_u = 3 \pm 1 \text{ MeV}$, $\langle \bar{d}d \rangle - \langle \bar{u}u \rangle = -(2.5 \pm 1) \cdot 10^{-3} \langle \bar{u}u \rangle$ (at a standard normalization point, $\mu = 0.5 \text{ GeV}$).

[†]Permanent address: Institute of Theoretical and Experimental Physics, Moscow 117259, Russia.

The QCD sum rules invented more than a decade ago is now well known to be a very useful tool to study properties of hadrons at intermediate energies and to get information on the basic parameters of QCD, such as quark masses and non-perturbative condensates. One of the problems addressed already in the pioneering papers[1] was the relation between the isotopic symmetry violation on the level of hadrons and the difference between u - and d -quark masses and condensates. It was shown that the observed $\rho - \omega$ mixing implies $(m_d - m_u)/(m_u + m_d) \sim 0.3$ and $\gamma \sim -1.5 \cdot 10^{-2}$, where

$$\gamma = \langle \bar{d}d \rangle / \langle \bar{u}u \rangle - 1 \quad (1)$$

ruling out a solution with $m_u = 0$, $m_d \neq 0$. The quark mass difference was estimated before the advent of QCD sum rules in refs.[2, 3] with the result $m_d - m_u \approx 3 \text{ MeV}$ and $m_d + m_u \approx 11 \text{ MeV}$. The difference of condensates was later estimated within the QCD sum rule framework in a number of papers[4] with the result $\gamma = -(3 \div 10) \cdot 10^{-3}$. The difference of condensates was also obtained in the framework of chiral perturbation theory[5], which gives $\gamma \approx -8 \cdot 10^{-2}$, provided $m_d - m_u = 3 \text{ MeV}$, $1 - \langle \bar{s}s \rangle / \langle \bar{u}u \rangle \approx 0.2$ and $m_s \approx 150 \text{ MeV}$. Recently isospin violation in QCD sum rules for the nucleon, Σ and Ξ was considered[6] and the following results were obtained: $m_d - m_u = 3 \pm 1 \text{ MeV}$, $\gamma = -(2 \pm 1) \cdot 10^{-3}$. Thus, while most predictions for $m_d - m_u$ agree and are grouped around 3 MeV , predictions for $\langle \bar{d}d \rangle - \langle \bar{u}u \rangle$ are more diverse and range within an order of magnitude. Moreover, arguments were given in ref.[7] that m_u may be equal to zero due to instanton contributions to the renormalization of the quark mass. Thus, additional independent estimates of differences between u - and d -quark masses and condensates are certainly welcome.

In this paper we will consider QCD sum rules for isospin mass splittings of D^* and D mesons and make use of the recently reported[8] new results on these splittings,

$$\begin{aligned} m_{D^{*+}} - m_{D^{*0}} &= 3.32 \pm 0.08 \pm 0.05 \text{ MeV} \\ m_{D^+} - m_{D^0} &= 4.80 \pm 0.10 \pm 0.06 \text{ MeV} \end{aligned} \quad (2)$$

to obtain such estimates. The sum rules are similar to those which were used in ref.[9] to successfully predict the mass splittings $m_{D_s^*} - m_{D^*} = 110 \pm 20 \text{ MeV}$ and $m_{D_s} - m_D = 120 \pm 20 \text{ MeV}$.

Let us start with the correlator of two pseudoscalar currents with quantum numbers of D , $j_5 = \bar{c}\gamma_5 q$, where q is either u , or d , at Euclidean momentum $-q^2 \gtrsim 1 \text{ GeV}^2$,

$$C^q = i \int d^4x e^{iqx} \langle 0 | T \{ j_5(x), j_5^+(0) | 0 \rangle \quad (3)$$

and consider its variation δC^q as the light quark mass rises from zero to its actual value m_q . To estimate C^q it is possible to take into account only the unit operator and the quark condensate (Fig.1) in the operator product expansion, since the contribution of operators of higher dimension to the heavy-light correlators is negligible[10]. Using the expansion of the quark condensate in the quark mass

$$\langle q_\alpha^a(x) \bar{q}_\beta^b(0) \rangle = \left(-\frac{1}{12} \delta_{\alpha\beta} \langle \bar{q}q \rangle + \frac{i}{48} m_q \hat{x}_{\alpha\beta} \langle \bar{q}q \rangle \right) \delta^{ab} \quad (4)$$

where $\langle \bar{q}q \rangle$ itself also depends on m_q , it is easy to see that δC^q is a function of m_q and $\langle \bar{q}q \rangle - \langle \bar{q}q \rangle_0$ where the subscript 0 denotes the chiral limit. On the other hand, saturating the correlators in the standard manner by the corresponding lowest mass resonances D^+ and D^0 , subtracting the continuum from the contribution of the unit operator and applying the Borel transformation[1], $(s + Q^2)^{-1} \rightarrow M^{-2} \exp(-s/M^2)$, we arrive at the following sum rule for $\delta C^d - \delta C^u$

$$\begin{aligned} -\frac{\beta_{D^+} - \beta_{D^0}}{\beta_D m_D} M^2 + (m_{D^+} - m_{D^0})_{hadr} &= \frac{M^4 e^{m_D^2/M^2}}{2\beta_D^2 m_D} \\ \{ (m_d - m_u) [\frac{3m_c}{4\pi^2} (e^{-x} - e^{-y} - x(E_1(x) - E_1(y))) L^{4/9} - \frac{\langle \bar{q}q \rangle}{2M^2} e^{-x} (1 + \frac{m_c^2}{M^2})] \\ + \frac{\langle \bar{d}d \rangle - \langle \bar{u}u \rangle}{M^2} m_c e^{-x} L^{4/9} - (s_{D^+} - s_{D^0}) \frac{3}{8\pi^2 M^2 s_D} (s_D - m_c^2)^2 e^{-y} \} \end{aligned} \quad (5)$$

Here $E_1(x) = \int_x^\infty dt t^{-1} e^{-t}$, m_c is the charmed quark mass, s_D is the continuum threshold, $x = m_c^2/M^2$, $y = s_D/M^2$ and the residue of D meson into the current j_5 is $\beta_D^2 = f_D^2 m_D^4/m_c^2$, where f_D is the semileptonic decay constant defined by $\langle D | \bar{c} \gamma_\mu \gamma_5 q | 0 \rangle = -i f_D p_\mu$. The dependence of light quark masses and condensates on the normalization point μ in the operator product expansion is given by powers of $L = \ln(M/\Lambda)/\ln(\mu/\Lambda)$ where $\Lambda = 150 \text{ MeV}$ and we take $\mu = 0.5 \text{ GeV}$ which corresponds to $\langle \bar{q}q \rangle = -(0.24 \text{ GeV})^3$. Here $\beta_{D^+} = \beta_D + \delta\beta_D^d$, $\beta_{D^0} = \beta_D + \delta\beta_D^u$ and $s_{D^+} = s_D + \delta s_D^d$, $s_{D^0} = s_D + \delta s_D^u$ where β_D and s_D are the residue and the continuum thresholds in the chiral limit and $\delta\beta_D^{u,d}$ and $\delta s_D^{u,d}$ are the deviations of residues and of continuum thresholds from their values in the chiral limit.

Similar sum rules can be written for the correlator of two vector currents, $j_\mu = \bar{c} \gamma_\mu q$,

$$C_{\mu\nu}^q = i \int d^4x e^{iqx} \langle 0 | T \{ j_\mu(x), j_\nu^+(0) | 0 \rangle \quad (6)$$

In the case of the tensor structure $q_\mu q_\nu$ for which the sum rule is known to work better[11], we obtain

$$\begin{aligned}
& -\left(\frac{\beta_{D^{*+}} - \beta_{D^{*0}}}{\beta_{D^*} m_{D^*}} - \frac{(m_{D^{*+}} - m_{D^{*0}})_{\text{hadr}}}{m_{D^*}^2}\right) M^2 + (m_{D^{*+}} - m_{D^{*0}})_{\text{hadr}} = \\
& -\frac{m_{D^*} e^{m_{D^*}^2/M^2}}{2\beta_{D^*}^2} [(m_d - m_u) \langle \bar{q}q \rangle e^{-x} + (s_{D^{*+}} - s_{D^{*0}}) \frac{e^{-y} M^2}{4\pi^2} (1 - \frac{3m_c^4}{s_{D^*}^2} + \frac{2m_c^6}{s_{D^*}^3})] \quad (7)
\end{aligned}$$

It is important to emphasize that since we do not take into account perturbative two-loop diagrams with one photon exchange in the "theoretical" part of the sum rules, only the hadronic parts of the isospin splittings, $(\Delta m_D)_{\text{hadr}}$ and $(\Delta m_{D^*})_{\text{hadr}}$, enter eqs.(5) and (7). To obtain them we use a quark model estimate[12] for the photon cloud part of the mass difference $m_{D^+} - m_{D^0} = 1.7 \pm 0.5 \text{ MeV}$ and also take into account the electromagnetic hyperfine splitting

$$\delta m = -\frac{2\pi Q_c Q_q |\Psi(0)|^2}{3m_c m_q} [2S(S+1) - 3] \quad (8)$$

where S is the spin of the meson. The quark masses here are the constituent ones, $m_c \sim 1.7 \text{ GeV}$, $m_q \sim 0.3 \text{ GeV}$. The quark-antiquark wave function at the origin may be estimated using the relation $f_D^2 = 12|\Psi(0)|^2/m_D$. Using the estimate[10] $f_D = 170 \text{ MeV}$, we get that the hyperfine electromagnetic interaction contributes $\sim 0.5 \text{ MeV}$ to $D^+ - D^0$ and $\sim -0.17 \text{ MeV}$ to $D^{*+} - D^{*0}$ mass splittings. Using the full experimental mass differences from eq.(2) we thus have the mass splittings to be used in the sum rules in eqs.(5) and (7)

$$\begin{aligned}
(m_{D^{*+}} - m_{D^{*0}})_{\text{hadr}} &= 1.8 \pm 0.5 \text{ MeV} \\
(m_{D^+} - m_{D^0})_{\text{hadr}} &= 2.6 \pm 0.5 \text{ MeV} \quad (9)
\end{aligned}$$

From the sum rules in eqs.(5) and (7) it follows that in the working region for the Borel parameter M^2 , the left hand sides should be close to linear functions in M^2 whose extrapolations to $M^2 = 0$ give the corresponding hadronic mass splittings while the slopes give information on the residue differences, $\beta_{D^+} - \beta_{D^0}$ and $\beta_{D^{*+}} - \beta_{D^{*0}}$. To numerically analyze the sum rules we take $m_D = 1.87 \text{ GeV}$, $m_{D^*} = 2.01 \text{ GeV}$, the standard value of the quark condensate, $\langle \bar{q}q \rangle = -(0.24)^3 \text{ GeV}^3$, and $m_c = 1.35 \text{ GeV}$. For the residues and continuum thresholds we take the estimates obtained with sum rules in refs.[10, 11], $\beta_D^2 \approx \beta_{D^*}^2 \approx 0.2 \text{ GeV}^4$ and $s_D \approx s_{D^*} \approx 6 \text{ GeV}^4$.

Thus, $m_d - m_u$, $\langle \bar{d}d \rangle - \langle \bar{u}u \rangle$, the Borel parameter M^2 and the differences of continuum thresholds, $s_{D^+} - s_{D^0}$ and $s_{D^{*+}} - s_{D^{*0}}$ are the fitting parameters of the sum rules in eqs.(5) and (7). The working regions in M^2 determined by the requirement of controllable contributions

from non-perturbative power corrections and continuum in the chiral limit for u - and d -quarks are [10, 11] $1 \text{ GeV}^2 \lesssim M^2 \lesssim 2 \text{ GeV}^2$ for the pseudoscalar channel and $1.5 \text{ GeV}^2 \lesssim M^2 \lesssim 2.5 \text{ GeV}^2$ for the vector channel. We will let the difference of thresholds vary around an estimate $s_{D^+} - s_{D^0} \sim s_{D^{*+}} - s_{D^{*0}} \sim (m_d - m_u)\sqrt{s_D}$ which implies that the threshold and the quark mass differences are of the same sign.

In Fig.2 we show the result of the sum rule calculation of the D^* mass difference by plotting $f_V(M^2) - M^2 df_V(M^2)/dM^2$, where $f_V(M^2)$ is the r.h.s. of eq.(7). From eq.(7) one sees that only the quark mass difference enters this sum rule and one might hope to fix $m_d - m_u$ from it. The term proportional to $s_{D^{*+}} - s_{D^{*0}}$ has the same sign as the term proportional to $m_d - m_u$ and in the whole working interval in M^2 contributes less than 50% to $f(M^2) - M^2 df(M^2)/dM^2$ at $m_d - m_u = 3 \text{ MeV}$. The values $m_d - m_u \gtrsim 4 \text{ MeV}$ cannot be made compatible with $(m_{D^{*+}} - m_{D^{*0}})_{\text{had}}$ from eq.(9) at any $s_{D^{*+}} - s_{D^{*0}} > 0$. At $m_d - m_u \lesssim 4 \text{ MeV}$ the compatibility can be achieved by varying $s_{D^{*+}} - s_{D^{*0}}$. However, at $m_d - m_u \lesssim 2 \text{ MeV}$ the continuum term is dominant in the whole working region in M^2 and the sum rule is not reliable.

As for the D mass difference, the corresponding sum rule in eq.(5) is contributed both by $m_d - m_u$ and $\langle \bar{d}d \rangle - \langle \bar{u}u \rangle$. In Fig.3 we show the numerical results for $m_d - m_u = 3 \text{ MeV}$ and two different values of $\gamma = \langle \bar{d}d \rangle / \langle \bar{u}u \rangle - 1$. One can see that the value $\gamma = -2.5 \cdot 10^{-3}$ is consistent with the $(m_{D^+} - m_{D^0})_{\text{had}}$ from eq.(9), while a higher value $\gamma = -6 \cdot 10^{-3}$ is definitely excluded.

In summary, we conclude from our numerical analysis of the sum rules for isospin mass splittings in D and D^* mesons that $m_d - m_u = 3 \pm 1 \text{ MeV}$ and $\gamma = -(2.5 \pm 1) \cdot 10^{-3}$. The result for $m_d - m_u$ is consistent with the earlier estimates and together with the relation $(m_u + m_d)\langle \bar{u}u \rangle = -F_\pi^2 m_\pi^2$ excludes the option $m_u = 0$ advocated in ref.[7]. Our estimate for the condensate difference parameter γ supports the value obtained in ref.[6] from the isospin splittings in baryons and differs from results of earlier papers[4, 5]¹.

We are grateful to H. Leutwyler for useful discussions. One of the authors (V.L.E.) acknowledges the warm hospitality extended to him at the Universities of Minnesota and Bern. This work was supported in part by the US Department of Energy under contract DE-FG02-87ER40328 and by Schweizerischer Nationalfonds.

¹For a discussion of possible reasons for this disagreement, see ref.[6].

References

- [1] M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, Nucl. Phys. B **147**, 385, 448, 519 (1979).
- [2] J. Gasser and H. Leutwyler, Nucl. Phys. B **94**, 269 (1975).
- [3] S. Weinberg, in: *Festschrift for I.I. Rabi*, ed. L. Motz, NY Academy of Sciences, New York, 1978.
- [4] P. Pascual and R. Tarrach, Phys. Lett. B **116**, 443 (1982);
E. Bagan, A. Bramon, S. Narison and N. Paver, Phys. Lett B **135**, 463 (1984);
C.A. Dominguez and M. Loewe, Phys. Rev. D **31**, 2930 (1985);
C.A. Dominguez and M. Loewe, Ann. Phys. (NY) **174**, 372 (1987);
S. Narison, Rev. Nuovo Cim. **10**, 1 (1987); *QCD Spectral Sum Rules*, World Scientific
Lecture Notes in Physics, v.26, Singapore, 1989;
E.G. Drukarev, St.-Petersburg Nuclear Physics Institute Report, PNPI-1780 (1992).
- [5] J. Gasser and H. Leutwyler, Nucl. Phys. B **250**, 465 (1985).
- [6] C. Adami, E.G. Drukarev and B.L. Ioffe, Marmal Aid Preprint Series In Theoretical
Nuclear Physics, MAP-151 (1993).
- [7] Choi, Phys. Lett. B **292**, 159 (1992);
Nucl. Phys. B **383**, 58 (1992).
- [8] D. Bortoletto et al, Phys. Rev. Lett. **69**, 2046 (1992).
- [9] B.Yu. Blok and V.L. Eletsky, Sov. J. Nucl. Phys. **42**, 787 (1985).
- [10] T.M. Aliev and V.L. Eletsky, Sov. J. Nucl. Phys. **38**, 936 (1983).
- [11] V.L. Eletsky and Ya.I. Kogan, Z.Phys. C **28**, 155 (1985).
- [12] J. Gasser and H. Leutwyler, Phys. Rep. **87**, 78 (1982).

Figure Captions:

- Fig. 1: The diagrams taken into account in the sum rules.
- Fig. 2: Sum rule for D^* mass splitting: $f_V(M^2) - M^2 df_V(M^2)/dM^2$ versus M^2 , where $f_V(M^2)$ is the r.h.s. of eq.(7) for $m_d - m_u = 1 \text{ MeV}$ (a), 3 MeV (b) and 5 MeV (c). In each of the three cases the lower and the upper curves correspond to $s_{D^{*+}} - s_{D^{*0}} = 0$ and 0.005 GeV^2 , respectively. The dotted horizontal lines are the boundaries set by eq.(9).
- Fig. 3: Sum rule for D mass splitting: $f_P(M^2) - M^2 df_P(M^2)/dM^2$ versus M^2 , where $f_P(M^2)$ is the r.h.s. of eq.(5) for $m_d - m_u = 3 \text{ MeV}$ and a) $\gamma = -2.5 \cdot 10^{-3}$; b) $\gamma = -6 \cdot 10^{-3}$. The numbers at the curves correspond to the value of $s_{D^+} - s_{D^0}$ (in GeV^2). The dotted horizontal lines are the boundaries set by eq.(9).